Investigation of Melia azedarach core and fruit adsorbent to remove Methylene blue from synthetic wastewater

Ali Asghar Najafpoor¹, Mojtaba Davoudi², Ali Akbar Dehghan³, Sepideh Bakhti^{*4}, Shiva Ghaderifar⁴, Elahe Ahmadi⁴

1- Associate Professor, Health Sciences Research Center, Department of Environment Health Engineering, School of Health, Mashhad University of Medical Science, Mashhad, Iran

2- Assistant Professor Department of Environmental Health Engineering, School of Health, Torbat Heydarieh University of Medical Sciences, Torbat Heydariyeh, Iran

3- PhD student of Environment Health Engineering, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

4- M.Sc student, Health Sciences Research Center ,Department of Environment Health Engineering, School of Health, Mashhad University of Medical Science, Mashhad, Iran

*Corresponding address: Health Sciences Research Center, Department of Environment Health Engineering, School of Health, Mashhad University of Medical Science, Mashhad, Iran. Tel: 09382240880 Email address:bakhtis921@mums.ac.ir

Abstract

Background & Aim: Dyes are one of the most important pollutants in textile wastewater that are toxic, carcinogenic, mutagenic and non-biodegradable. Melia azedarach core and fruit are among the inexpensive natural absorbents that have important place in the treatment of textile wastewater. This study was conducted to determine the efficiency of Melia azedarach core and fruit in removal of Methylene blue from synthetic wastewater.

Methods: In this study, structural characteristics of absorbent were analyzed using Fourier transform infrared spectroscopy (FTIR) and Scanning electron microscope (SEM). Also the effect of various parameters such as initial dye concentrations (10-100 mg/l), adsorbent dosage (0.1-1.1 g/l), contact time (0-60 min) and solution pH (4-10) on the removal of dye was investigated and at the end, kinetics and adsorption isotherm were also determined. Dye concentration was analyzed by UV/VIS spectrophotometer at wavelength of 665 nm.

Results: Maximum removal of dye (85%) was observed in dye concentration of 32.5 mg/l, adsorbent dose of 0.85 g/l, detention time of 45 min and pH=8.5. Initial concentration had the greatest impact on color removal (P=0.0001). The results showed that data were explained acceptably by pseudo second-order model (R^2 =0.95_0.99) and Langmuir isotherm model (R^2 =0.98).

Conclusion: Melia azedarach core and fruit have the ability to remove methylene blue. Since in this study, the optimum pH was obtained in alkaline range and most textile wastewaters are in this range, it is an appropriate and cost-effective adsorbent to remove dye from the wastewater of textile industries.

Keywords: Adsorption; Melia azedarach; Methylene blue; Wastewater textile

Page 29