Assessing the effect of drug tolerance due to chronic administration of morphine and salicylate on synaptic plasticity

Masoumeh Gholami^{1*}, Mehdi Sadegh²

1. Neurosciences Research Center, Torbat Heydariyeh University of Medical Sciences, Torbat Heydariyeh, Iran

2. Department of Physiology, Faculty of Medicine, Arak University of Medical Sciences, Arak, Iran.

*Corresponding Address: Basic Sciences and Neurosciences Research Center, Torbat Heydariyeh University of

Medical Sciences, Torbat Heydariyeh, Iran

Email address: Gholamim3@thums.ac.ir

Abstract

Background & Aim: Salicylates and opioids are widely used in chronic pain relief. Chronic use

of these drugs reorganizes synaptic function, especially experience-dependent plasticity in brain

regions. Therefore, in this study the effects of chronic administration of salicylate and morphine

on synaptic plasticity were investigated.

Methods: in this review, Elsevier, Science Direct, PubMed and Google Scholar databases were

searched for the following keywords: salicylate, morphine, drug tolerance, and synaptic

plasticity. At the end, 62 of the obtained reports were included in the study.

Results: In addition to induction of tolerance to anti-nociceptive effect and cross-tolerance,

chronic salicylate administration as like as morphine has long-lasting effects on hippocampal

neuronal networks which are manifested as excitability of neurons and ability of activity and

experience-dependent plasticity. Some of these effects can be documented in the hippocampus-

related functions such as spatial learning and memory.

Conclusion: Considering the effects of salicylate and morphine on the nervous system and

synaptic transmission, the effects of these drugs on the processing of input data and as a result on

cognitive functioning would not be unlikely, thus necessitating further behavioral and

electrophysiological studies.

Keywords: Salicylate, Morphine, Drug tolerance, Synaptic plasticity